


**Europäisches Patentamt**  
**European Patent Office**  
**Office européen des brevets**

Publication number:

**0 147 483**  
**A1**

**EUROPEAN PATENT APPLICATION**

Application number: 83113183.4

Int. Cl.<sup>4</sup>: **A 23 G 9/02**  
**A 23 G 9/04**

Date of filing: 28.12.83

Date of publication of application:  
10.07.85 Bulletin 85/28

Designated Contracting States:  
AT BE CH DE FR GB IT LI LU NL SE

Applicant: **THE PILLSBURY COMPANY**  
200 South 6th Street  
Minneapolis Minnesota 55402(US)

Inventor: **Desai, Nitin**  
872 Raymond Apt. 10  
St. Paul Minnesota 55114(US)

Inventor: **Delaney, Rory A.M.**  
16708 Picha Lane  
Minnetonka Minnesota 55343(US)

Inventor: **Brouwer, Peter**  
28th Avenue North  
Plymouth Minnesota 55441(US)

Inventor: **Huang, Victor T.**  
3813 72nd Avenue North  
Brooklyn Center Minnesota 55429(US)

Representative: **Patentanwälte Grünecker, Dr.**  
**Kinkeldey, Dr. Stockmair, Dr. Schumann, Jakob, Dr.**  
**Bezold, Melster, Hilgers, Dr. Meyer-Plath**  
Maximilianstrasse 58  
D-8000 München 22(DE)

**Stable aerated frozen food product.**

A process for preparing an aerated frozen food product such as mellorine or a shake beverage wherein water, fat, protein, emulsifier and stabilizer are blended together to form a mix which is homogenized to form an oil-in-water emulsion and is subsequently whipped under freezing conditions. Improved stability is achieved in the product by selecting a specific oil having high solid fat content at room temperature, homogenizing the mix of ingredients under conditions sufficient to form an emulsion having a relatively narrow distribution of small diameter fat globules, and aging the emulsion to crystallize the fat globules prior to whipping. Stable aerated frozen products provided in accordance with this invention can be stored and distributed at 0°F without loss of quality and have a variety of commercial uses.

EP 0 147 483 A1

5

10

STABLE AERATED FROZEN FOOD PRODUCT

Frozen aerated food products to which this invention  
15 is directed include frozen desserts such as ice cream, ice  
milk, mellorine, sherbet, frozen custard, frozen pudding  
and the like, frozen shake beverages such as milk shakes,  
and frozen shake concentrate. The popularity of these  
products is attributable to their convenience, widespread  
20 availability, nutritive value and appealing forms, colors  
and flavors. In part due to the familiarity of such  
frozen products to the consumer, to be acceptable they  
must not only have suitable flavor and appearance, but  
also possess a complex set of physical properties which  
25 provide the characteristics which consumers have come to  
expect. In addition, the nature and composition of the  
most popular aerated frozen products are set by various  
standards promulgated by the U.S. Food and Drug Adminis-  
tration as well as the individual states. Such rigid  
30 consumer acceptance criteria, regulatory standards and the  
complex nature of the interactions which produce a  
satisfactory product make it extremely difficult to  
reformulate such aerated frozen products in an attempt to  
provide improvements. Solutions to one set of problems  
35 quite often generate other problems which make the product  
unacceptable.

Stability is one major problem of conventional aerated frozen products. For example, ice cream and related aerated frozen desserts are stored, distributed, delivered and sold at deep freeze temperatures, i.e. -20°F, to give some measure of stability to these products and protect against deterioration of their special textural properties. This storage and distribution system, however, has many drawbacks both from the consumer's and manufacturer's point of view. One problem is that, at deep freeze temperatures, the product is so hard that it is difficult to scoop and serve immediately from the deep freeze.

Another major problem is that temperature fluctuations invariably occur during storage and handling of aerated frozen products which result in the thawing and refreezing of the product, seriously deteriorating its quality over time. This "heat shock" is characterized by the formation of large ice crystals, and results in excessive hardening and gritty mouthfeel. Heat shock frequently also produces a separation of the syrupy aqueous phase from the air and fat matrix which ultimately make the appearance and texture of the product unacceptable.

25

Several solutions to these problems have been attempted in the past. For example, increased stability has been imparted to aerated frozen desserts through the use of various stabilizers and stabilizer combinations. The drawback of this approach is that stabilizers frequently are required in such great quantities that the feel of the product in the mouth is altered, producing a cloying, gummy or greasy sensation. Another approach has been to increase the amount of added sugars relative to the fat and/or water in the product, but the large amount of sugar required to provide acceptable stability often

results in too sweet a taste or unacceptable crystal formation. Still another approach has been to increase the amount of incorporated air, i.e., "overrun", and thereby decrease the amount of freezable water per unit  
5 volume. This attempt at reformulation, however, not only results in a texture which is more characteristic of a whipped topping than of a frozen dessert like ice cream, but is limited by the applicable standards of identity relating to required weight per volume and solids content.

10

As a result of the instability of most aerated frozen desserts, producers have been required to manufacture the products locally to avoid the long periods of storage and temperature variations which are associated with national  
15 distribution from a few centrally located plants. Since most producers, distributors and retailers rely on deep freeze temperatures for preserving the quality of ice cream-type desserts during storage, distribution and sale, totally separate systems are rerquired for such products  
20 than are used for other frozen foods, which can be distributed at temperatures ranging from 0°F to 10°F. A very extensive network of frozen food storage, distribution and retailing facilities has been established nationwide, and therefore it can be seen that it would be  
25 extremely advantageous from an economic and efficiency standpoint if an ice cream-type aerated frozen dessert of acceptable quality and stability could also utilize this system.

30 Milkshakes and similar frozen shake beverages are representative of another type of aerated frozen food product in which poor stability has limited the storage and distribution systems which the manufacturer can utilize. As is well-known, milkshakes are usually  
35 manually prepared or dispensed from a commercial establishment for consumption on the premises. Various

attempts have been made in the past to provide products which are more widely available or are more susceptible to automated procedures. For example, milkshakes have simply been packaged in individual cups and frozen solid for  
5 distribution. The product is then warmed under controlled conditions to room temperature, agitated or otherwise mixed, and served. Another approach has been to prepare and distribute a frozen milkshake concentrate which serves as a base to which milk or water is added with agitation.  
10 Still another approach has been to aseptically can or bottle a prepared milkshake which is consumed after being chilled, shaken and opened. All of these prior approaches have either required unacceptably large amounts of time or space, required a reconstititional step which is either  
15 difficult or time-consuming and/or have provided milkshakes which do not have the desired creaminess, coldness and consistency of commercially prepared milkshakes.

20 Since shakes are extremely popular items at fast food restaurants, it is of course important to reduce the time and labor involved in preparing large quantities of shakes manually. The present invention provides stable shakes which can be prepared in individual containers in central  
25 locations and shipped in a frozen state to various restaurant locations where they are stored and served at refrigeration temperatures without any on-site preparation.

30 Accordingly, it is an object of this invention to provide aerated frozen food products which are stable for sufficient periods of time to allow national distribution under conditions and temperatures encountered in 0°F frozen food systems. Another object is to provide an  
35 aerated frozen product which attains the aforesaid stability without a basic reformulation which would bring

the product outside the limits of consumer acceptance and regulatory standards. A still further object of the invention is to provide aerated frozen dessert products having the above attributes which are also soft and scoopable at freezer temperatures. Another object is to provide shakes and similar beverages having the above attributes which retain the desirable characteristics of commercially prepared milkshakes when served at refrigeration temperatures (e.g. 15°F-25°F). A related object is to provide a frozen shake concentrate which can readily be reconstituted by the consumer into a high-quality shake.

In accordance with one aspect of the invention, a process for preparing an aerated frozen food product, wherein water, fat, protein, emulsifier and stabilizer are blended together to form a mix and said mix is homogenized to form an oil-in water emulsion which is subsequently whipped under freezing conditions, comprises:

selecting an edible oil which has an SFI of at least 25 to 70°F;

homogenizing said mix to form an emulsion of fat globules having a particle size wherein the average  $d_{vs}$  value is 0.2 to 1.5 microns, and a particle size distribution wherein the ratio of  $d_{vs}$  to  $d_{max}$  has a quotient in the range from 9 to 14;

aging the homogenized mix to crystallize said fat globules prior to said whipping and freezing; and

substantially retaining said particle size and particle size distribution after said whipping and freezing.

In accordance with the present invention, therefore, the fat component of the aerated frozen food product comprises an edible oil which has a solid fat index (SFI) of at least 25 at 70°F. The mix containing this fat is  
5 homogenized to form an emulsion of fat globules having an average volume/surface diameter ( $d_{vs}$ ) of 0.2 to 1.5 microns, and a ratio of maximum fat globule diameter ( $d_{max}$ ) to  $d_{vs}$  having a quotient of from 9 to 14. The emulsion is subsequently aged to crystallize the fat  
10 globules prior to whipping. The vast majority of the fat in the product, i.e. at least 94%, is in the form of fat dispersed in the aqueous phase with only a small amount, i.e. less than 6%, in the form of free or de-emulsified fat. The overrun of the aerated frozen food product is  
15 preferably from 50% to 110%.

The stable aerated frozen food products of the invention are extremely versatile and can be given a variety of textures and consistencies by varying the  
20 formulation. A preferred embodiment of the invention comprises a stable mellorine product having 52% to 57% water in the form of ice crystals and an aqueous phase, ~~10% to 15%~~ of the aforementioned edible oil, 4% to 15% non-fat milk solids, and minor but effective amounts of  
25 stabilizer and emulsifier. Preferably, the whipping step is sufficient to provide an overrun in the range of 60% to 90% in order to produce mellorine having the textural characteristics and appearance of conventional ice cream. Products having the characteristics of soft-serve ice  
30 cream at 0°F may further comprise a freezing point depressant, preferably in the form of added sugars or other soluble solutes.

Ready-to-serve shake beverages having improved  
35 stability are also provided in accordance with this invention by increasing the amount of water in the above-

mentioned formulation and lowering the overrun to achieve the desired shake consistency. Preferred shakes of the invention comprise about 63% to 68% water, 3% to 9% of the aforementioned edible oil, and about 50% to 70% overrun.

- 5 These shake products can be distributed through conventional frozen food systems in solid frozen form and then stored for several days at refrigeration temperatures (15°F to 25°F) before consumption without losing the desirable characteristics of commercially prepared
- 10 milkshakes. The invention can also be employed to provide a frozen concentrate to which water or other liquids are added at room temperature with stirring to produce shake beverages.

- 15 It has quite unexpectedly been found that the selection and processing of the fat component during the preparation of the aforementioned aerated frozen food products are critical elements in achieving the unique stability of this invention. Because of the solid fat
- 20 index (SFI) and other properties of the fat used, the small dispersed fat globules are substantially all crystallized during the aging step and a significant amount of these solid crystals are retained when the product is exposed to room temperature. Without intending
- 25 to be bound by theory, it is thought that the mobility of the aqueous phase during thawing and refreezing of the product is significantly restricted by these finely distributed fat crystals, thereby resulting in products which develop less iciness and retain overrun better when
- 30 exposed to heat shock than otherwise could be provided. This property in turn is thought to account in large part for the improved shelf life stability of the present products at typical frozen food temperatures.



Preparation in accordance with this invention results in aerated frozen food products which have sufficient stability to retain their desirable organoleptic properties for at least 50 days, and preferably as long as 5 180 days, at conditions which are normally encountered in typical 0°F frozen food storage, distribution and retailing systems. As a result, the products may be manufactured in a relatively few plants and distributed nationally and/or internationally using the same 10 facilities as other frozen food items such as frozen vegetables, frozen entrees and frozen pizza. Frozen dessert products of the invention do not require a separate deep freezer in the retail store, and can be sold and/or consumed directly from the frozen food case or home 15 freezer. The shake beverages of the invention can be delivered to the restaurant site frozen solid in individual containers and then stored and dispensed directly from a dispenser having a chamber in which the product has equilibrated to a temperature from 15°F to 20 25°F. Alternately, shakes can easily be prepared at home from a frozen concentrate of the invention stored in the home freezer. Preferably, 2 or 3 parts of frozen concentrate are diluted at room temperature (65°F to 75°F) with one part of water, or some other suitable liquid such 25 as milk or juice, and then the diluted concentrate is stirred with a spoon to form the shake.

As briefly described above, the oils suitable for use in the practice of this invention have a Solid Fat Index 30 (SFI) at 70°F of at least 25. SFI values are readily determined in accordance with standard analytical procedures using a dilatometer. Although the oil has a significant solid fat content at 70°F, it should be liquid at that temperature or easily liquefiable on warming and 35 be substantially completely melted at about 100°F or slightly above. As used herein and in the claims the term

"oil" is intended to designate liquid fats meeting these requirements. Generally, the preferred oils are hydrogenated vegetable oils which have a (capillary) melting point between 70°F and 106°F, or blends of such vegetable oils. Suitable oils are derived from coconut, soybean, cottonseed, corn, palm kernel, peanut or the like. The solid fat in the most preferred oils melts quickly between 70°F and 100°F and these oils exhibit a sharp peak on a Differential Scanning Calorimetry curve. Since ice cream, ice milk, and milkshake standards require the use of butter fat, it is contemplated that high melting point fractions of butter oil meeting the SFI requirements of the invention could also be used to produce stable ice cream, ice milk, and milkshake products in accordance with this invention.

The amount of the fat component will vary depending on the aerated frozen food product being prepared and the particular organoleptic properties desired. Generally, the amount of the fat component comprises about 3% to 30% by weight of the frozen product, with the preferred mellorine products comprising 5% to 30% fat, most preferably 10% to 15% fat. The most preferred shake products of the invention comprise from 3% to 9% fat.

The fat component of this invention can be mixed with other ingredients at ambient temperatures but is preferably injected at an elevated temperature, e.g. about 160°F, into the aqueous mix containing the other ingredients. The resultant mixture is then homogenized, primarily to reduce the size of the fat globules and form a stable oil-in-water emulsion. In accordance with the preferred embodiment of the invention, the mix is homogenized at a low pressure, i.e., a total pressure below 6000 psig, to provide a uniform distribution of small fat globules in the emulsion.

The designations " $d_{vs}$ " and " $d_{max}$ " used herein and in the appended claims to characterize the particle size of the emulsion are derived from the spectroturbidimetry method described in detail by Walstra [P. Walstra, 5 Estimating Globule-Size Distribution of Oil-in-Water Emulsions by Spectroturbidimetry. J. Coll. Interf. Sci. Vol. 27, No. 3, P. 493-500 (1968); and P. Walstra, Light Scattering by Milk Fat Globules, neth. Milk and Dairy J., Vol. 19, No. 2, P. 93-109 (1965)]. In general, this 10 technique is based on the principle that the turbidity of an emulsion such as milk can be used to determine the average particle size and particle size distribution of the dispersed oil phase. In this method, a light beam is directed through the emulsion and the light scattering 15 coefficient is taken as a function of a dimensionless number derived from the particle size of the suspended fat globules and the wavelength of the light. The transmittance and absorbance of the light beam incident on the emulsion contained in a standard cell is measured for 20 various wavelengths using a spectrophotometer, and then these values are used to construct a light scattering spectrum. Homogenized emulsions typically show a log normal frequency-volume distribution, so a computer is used to generate a theoretical light scattering spectrum 25 based on a log normal particle size distribution. The volume/surface average diameter of the fat globules (herein and in the claims referred to by the designation  $d_{vs}$ ) and the maximum emulsion diameter (herein and in the claims referred to by the designation  $d_{max}$ ) are mathe- 30 matically calculated from the "fit" of the experimentally derived curve to the theoretical curve.

In accordance with the present invention, the mix is homogenized to provide a  $d_{vs}$  of 0.2 to 1.5 microns, 35 preferably 0.2 to 0.5 microns, with the ratio of  $d_{max}/d_{vs}$  having a quotient of from 9 to 14. As previously stated,

this relatively narrow distribution of small fat globules in combination with the solid fat characteristics of the oil used has been found to be very important in achieving the improved shelf-life characteristics of the present invention.

Homogenizers commonly used in ice cream manufacture may be employed to homogenize the mix. Although homogenizing techniques not utilizing pressure may be used, a homogenization pressure not in excess of a total of 6000 psig is preferred. In fact, one advantage of the present invention is that the desired homogenization can be achieved at pressures lower than that usually employed in conventional ice cream manufacture, e.g. pressures below 3000 psig. Homogenization is preferably accomplished at temperatures of 110°F to 180°F.

The vigorous agitation of the mix during freezing and whipping inevitably results in some destabilization of the fat emulsion in the form of fat "churnout" or "de-emulsification", i.e., a coalescence wherein the fat globule loses its identity as a dispersed entity and forms pools, and/or "clumps" together to form larger particles. In addition, the emulsion prior to whipping may inherently contain some of this "free" fat. The current understanding in the ice cream art is that such physical properties as dryness and stiffness are related to the degree of de-emulsification of the fat. It is believed that too little de-emulsification results in a wetter-appearing and less stiff ice cream, whereas too much de-emulsification may result in a courser texture. In addition, a coating of de-emulsified fat around the air cells tends to improve the foam stability of the product and is considered desirable. However, too much de-emulsified fat in the continuous phase may give a fatty or too creamy sensation to the frozen product.

It has quite surprisingly been found that the products of this invention exhibit considerably less fat de-emulsification, or free fat, than conventional frozen desserts and shake products. In the present products, at least 94% of the total fat is in the emulsified form and less than 6% of the total fat is in the de-emulsified form, whereas conventional ice cream, for example, typically has about 13% of the total fat in the de-emulsified form. The relatively narrow distribution of small fat crystals produced as a result of the homogenization and aging steps of the invention is substantially retained after the freezing and aeration step and this in turn is believed to contribute to this low incidence of de-emulsified fat.

15

De-emulsified fat can be extracted by hydrophobic organic solvents such as chloroform because of its "free" character, whereas emulsified fat cannot. Therefore, the level of de-emulsified fat in the frozen product can readily be determined by mixing a sample with chloroform, removing the chloroform soluble fraction, evaporating the chloroform and weighing the residue. De-emulsified fat so determined is then expressed as a percentage of the total fat.

25

The fat component of the invention may be used with mono and diglyceride emulsifiers and other emulsifiers normally used in aerated frozen food products and/or permitted by the applicable standards of identity. Such emulsifiers are typically used in amounts ranging from 0.1% to 0.5%, preferably less than 0.3%. The emulsifiers may be conveniently melted with the fat component and metered together into the mix prior to homogenization.

30

The mix may be pasteurized before or after homogenization in accordance with known commercial procedures. For example, continuous high temperature/short time (HTST) pasteurization is the most common method used by the  
5 larger ice cream plants and is suitable for the practice of the present invention. Minimum pasteurization standards for ice cream mix recommended by the U.S. Public Health Service are 175°F for 25 seconds for the HTST method, and 155°F for 30 minutes for the alternative  
10 holding method.

Following homogenization and pasteurization, the mix is rapidly cooled and held ("aged") at 30°F to 45°F for a time sufficient to cause extensive crystallization of the  
15 fat globules, usually from 2 to 12 hours. It is believed that the resultant distribution of small fat crystals which retain a significant amount of solid fat at room temperature not only aids in the freeze-thaw stability of the product by providing physical barriers limiting the  
20 mobility of the aqueous phase, but also contributes to improved physical properties of the frozen product, such as a smoother texture.

After sufficient aging, the mix is simultaneously  
25 aerated ("whipped") and frozen in the conventional manner. In a continuous freezer, the mix is rapidly cooled to freezing temperature and air is simultaneously incorporated to form the basic foam structure of small uniformly dispersed air cells. These processes take place  
30 under conditions of vigorous agitation provided by a "mutator", a rotating cylindrical device equipped with scraper blades and a whipping mechanism. The residence time during freezing is usually on the order of 25 to 30 seconds. As freezing proceeds, the mass becomes  
35 increasingly viscous and is finally extruded into containers from the freezer in a stiff plastic condition.

The product may then be further frozen (hardened) at a temperature of about -40°F to -80°F prior to storage at typical frozen conditions.

5       The amount of incorporated air, i.e., the "overrun",  
is important to the overall stability of aerated frozen  
products, yet must be controlled within certain limits to  
provide an acceptable texture and appearance. Generally,  
the higher the overrun, the lower the amount of water per  
10 unit of volume, and therefore the more resistant the  
product is to the formation of the large ice crystals  
after heat shock. Although high overrun aerated frozen  
products exhibit increased stability, increasing the  
overrun to improve the stability will not always produce  
15 an acceptable product since there are limits placed on the  
amount of air incorporated in the product, both by the  
various standards of identity and consumer preference.  
For example, federal regulations on solids content and  
weight/volume relationships for ice cream and related  
20 products, such as mellorine and ice milk, effectively  
limit the overrun to about 110%. Frozen desserts having  
overruns in excess of this amount tend to exhibit textural  
properties which are not characteristic of ice cream, and  
therefore are not preferred by consumers. The practice of  
25 the present invention is particularly advantageous for  
stabilizing ice cream-type frozen desserts without  
requiring overruns over 110%. Although products of this  
invention generally comprise less water than conventional  
ice cream-type products, they have similar overrun so as  
30 to retain the familiar textural properties of ice cream.  
Preferably, the frozen mellorine desserts and shake  
concentrates described herein comprise from 45% to 68%  
water, with 52% to 57% water being the most preferred  
range. Overrun for these products is usually from 50% to  
35 110%, preferably about 60% to 90%.

There are no existing standards which dictate the amount of overrun in milkshakes and similar frozen shake products. However, such products usually become too foamy to be acceptable at overruns of 80% or over, and too liquid or thin at overruns of 30% or below. For the ready-to-serve frozen shakes of this invention, the preferred water content is 63% to 68% and the preferred overrun is about 50% to 70%.

10 A stable foam (aerated) phase imparts a structural rigidity to the subject frozen food products and aids in limiting the mobility of the aqueous phase which causes the separation of the phases during prolonged storage. To this end, foam stabilizers are useful in the practice of  
15 the present invention. Among the most useful stabilizers are the hydrophilic colloids, or hydrocolloids, commonly referred to as "gums". These long-chain high-molecular-weight polymers disperse in the aqueous phase and/or interact with other product ingredients to provide a  
20 thickening or gelling effect which stabilizes the foam structure. They also contribute to the stability of the product by serving as water-binding ingredients which limit the mobility of the aqueous phase. Common gums suitable for use in the present invention include natural  
25 gums such as carrageenan, guar gum, locust bean gum, xanthan gum, gelatin, alginates, pectin, dextran, glucan and the like, as well as modified natural gums such as carboxymethyl cellulose, methylcellulose ether and other modified cellulose derivatives, modified starch,  
30 polyacrylic acid, and the like, and various mixtures thereof. The permitted quantities of gums may be governed by federal and/or state regulations, as well as by consumer preference regarding such physical properties as mouth feel and melt-down rate. The unique stability  
35 characteristics of the products of this invention allow



very small amounts of stabilizers to be used, preferably quantities much less than 1% and most preferably less than 0.3%.

5           Aerated frozen dessert products of the invention may be given hardness and texture at 0°F which simulate that of commercially available soft-serve ice cream. Conventional soft-serve ice cream products are usually prepared using machinery located on the premises where they are  
10 consumed and are served at temperatures slightly below freezing, e.g., 15°-25°F. Accordingly, these products characteristically have poor keeping qualities which prevent their effective distribution and storage off premises. If exposed to the deep freeze temperatures  
15 necessary to prolong their shelf life, conventional soft-serve products become very hard and cannot be scooped from a container for serving. To solve this problem, aerated frozen dessert products sometimes include greater than normal quantities of sugars, alcohols and other low  
20 molecular weight compounds to lower the freezing point of the aqueous phase to an extent where the product is soft and scoopable and has a soft-serve texture at freezer temperatures. Frozen shake concentrates are also made more easily dispersible by the addition of such compounds.  
25 It has been found that the use of such freezing point depressants is compatible with the practice of the present invention and that the unique stability of the products can readily be achieved with formulations including such compounds.

30

          The most useful freezing point depressants in the practice of this invention are sugars which not only have a sweetening effect, but also enhance the creamy texture of the product. The choice of sugars employed is  
35 controlled by the degree of freezing point depression desired and also by flavor and texture effects resulting

0147483

from the addition of such sugars. An insufficient amount of sugar will produce an off-taste in the product, whereas too much sugar will produce excessive sweetness and may mask other desirable flavors. Different sugars also lower  
5 the freezing point of the product to varying degrees, depending on their molecular weights.

Preferably, the amount of added sugars in the products of this invention is 18% to 33% by weight. It  
10 has been found that the most desirable balance of sweetness and textural properties may be achieved by employing sucrose in combination with one or more sugars which are less sweet; for example, sucrose may be combined with dextrose, lactose, low-conversion corn syrup, etc; in  
15 order to obtain the benefits of a relatively large proportion of sugar at a level of sweetness which is less than that resulting from the use of an equal amount of sucrose alone.

20 The products of this invention comprise 4% to 15% non-fat milk solids. Although non-fat milk solids are the preferred source of protein, it is contemplated that other suitable frozen dessert proteins, such as casein, caseinates, whey protein concentrate, egg protein, and  
25 soy, peanut and/or vegetable proteins may be used. Non-fat milk solids are the solids of skim milk and can be added in dry form or as condensed skim milk. These solids include proteins, minerals and milk sugar. Milk sugar adds to the sweet taste of the frozen product and proteins  
30 aid in the development of the desired overrun as well as contribute to the desirable texture and body of the product. The use of larger amounts of non-fat milk solids increases the effect that whipping has on the frozen product and increases the viscosity and resistance to  
35 melting of the composition. They also tend to lower the freezing point of the product. Non-fat milk solids which

contain not less than 2.5 mg/g undenatured whey protein nitrogen per gram of milk solids, as determined by the procedure set forth in American Dry Milk Institute Bulletin 916, are preferred.

5

The following examples are intended to illustrate the present invention, and are not to be construed as limiting the invention in any way:

10

Example I

A series of six frozen aerated desserts were prepared containing the following ingredients in parts by weight:

15

Parts/Weight

	Non-fat Dry Milk Solids	6.60
	Fat Component	12.00
	Dextrose	21.2
20	Sucrose	4.10
	Corn Syrup	2.40
	Xanthan Gum	0.264
	Carrageenan	0.004
	Emulsifier	0.10
25	Flavoring	0.40
	Coloring	0.84
	Water	52.10

The samples were made according to the following  
30 procedure:

48 pounds of the fat component was melted with 0.4 pounds of the emulsifier component by heating to a temperature of 110°F in a 70 gal. jacketed container to  
35 form the fat phase. 26.4 pounds of non-fat dry milk solids was dispersed in 206 pounds of water and the

0147483

remaining ingredients (122 pounds) were added to form the aqueous phase. The fat and aqueous phases were mixed together at 160°F, homogenized at a pressure of 1500 psi, and the emulsion thus formed was pasteurized at 175°F for 5 25 seconds. The mix was then cooled to between 30°F and 40°F and aged for 12 hours at 40°F. The aged mix was aerated in an ice cream freezer at about 16°F to provide an overrun of about 75%, extruded into suitable containers, hardened at a temperature of -40°F to -80°F and 10 stored in a 0°F freezer.

The SFI values of the fat components employed in the six samples were as follows:

15	Sample	<u>SFI at 100°F</u>	<u>Fat Component</u>	<u>SFI at 70°F</u>
	A		Coconut Oil	46.10
	B		Coconut Oil	37.40
20	C		Coconut Oil	26.60
	D		Soy/Coconut Oil Blend	20.00
	E		Soybean Oil	16.90
	F		Anhydrous Butter	13.80

25 The average fat particle size ( $d_{vs}$ ) of each sample was from 0.2 to 0.5 microns, and the  $d_{max}/d_{vs}$  quotient was in the range from 9 to 14. The amount of free fat for each sample was 2% to 3% using the chloroform extraction method.

30

The samples were subjected to conditions which were representative of those encountered in a typical 0°F distribution system. Samples in their containers on a "mock" pallet were removed from the 0°F freezer, kept for

2-4 hours at room temperature (about 70°F), and then placed back in the 0°F freezer for 20-22 hours. This procedure was repeated 4 or 5 times.

- 5        After undergoing the above freeze-thaw cycles, the samples were submitted to test panels consisting of 10 to 15 people, who were asked to rate them for iciness and softness. Samples A, B, and C incorporating the present invention were judged by the panels to be acceptable,  
10        whereas the remaining samples not incorporating the invention were determined to be unacceptable.

Example II

- 15        A Sample G employing the coconut oil of Sample B was prepared and subjected to heat shock in accordance with the procedure described in Example I, except that Sample G was prepared in a low shear device (i.e. an Oakes mixer) to provide an average particle size ( $d_{vs}$ ) of 6.4 microns  
20        and a ratio of  $d_{max}$  to  $d_{vs}$  having a quotient of 2.0. Sample G was considered to be unacceptably unstable to heat shock and was judged to be icy and coarse by the evaluation panel.

25

Example III

A frozen shake product was prepared using the procedure of Example I and the following formula:

		<u>Parts/Weight</u>
	Non-fat Dry Milk Solids	10.000
	Coconut Oil	5.715
5	Dextrose	14.000
	Sucrose	6.000
	Xanthan Gum	0.175
	Carrageenan	0.060
	Emulsifier	0.100
10	Flavoring	0.400
	Coloring	0.400
	Water	63.150
	Overrun	60%
15	This product (Sample H) was extruded into a 14 fl.	
	oz. waxed paper cup and placed in a 20°F refrigerator	
	along with a prior art frozen shake (Sample I) made in	
	accordance with Arbuckle U.S. Pat. No. 3,479,187. Samples	
	H and I were taken from the refrigerator, and along with	
20	Sample J, a typical fast food restaurant shake, they were	
	allowed to sit at room temperature (72.4°F) for 40	
	minutes. Both Sample I and J exhibited symptoms of	
	instability after this time. Sample J developed a phase	
	separation band about 2 cm wide at the bottom, and large	
25	(about 0.3 cm diameter) air bubbles on the top to give a	
	very foamy appearance. Sample I developed a rim of fluid	
	around an unappealing icy lump in the center of the shake.	
	However, there were no significant changes observed in	
	Sample H, during this time period, thereby demonstrating	
30	the superior stability of the shake beverages of the	
	present invention when compared to prior art products.	

0147483

CLAIMS:

1. A process for preparing an aerated frozen food product, wherein water, fat, protein, emulsifier and  
5 stabilizer are blended together to form a mix and said mix is homogenized to form an oil-in-water emulsion which is subsequently whipped under freezing conditions, characterized in that it comprises:

10 selecting an edible oil which has an SFI of at least 25 at 70°F;

homogenizing said mix to form an emulsion of fat globules having a particle size wherein the  
15 average  $d_{vs}$  value is 0.2 to 1.5 microns, and a particle size distribution wherein the ratio of  $d_{vs}$  to  $d_{max}$  has a quotient in the range from 9 to 14;

20 aging the homogenized mix to crystallize the fat globules prior to the whipping and freezing; and

25 substantially retaining the particle size and particle size distribution after the whipping and freezing.

2. A process as defined in claim 1, characterized in  
30 that the mix is homogenized at a pressure less than 6000 psig.

3. A process as defined in claim 1 or claim 2 characterized in that the homogenized mix is aged at 30°F to  
35 45°F for 2 to 12 hours prior to whipping and freezing.

4. A process as defined in any one of claims 1 to 3, characterized in that the aged mix is whipped to an overrun of 50% to 110% to provide a frozen dessert product.

5

5. A process as defined in any one of claims 1 to 3, characterized in that the aged mix is whipped to an overrun of about 50% to 70% to provide a frozen shake product.

10

6. A process for preparing a stable aerated frozen dessert product, characterized in that the process comprises the steps of:

15

mixing 45% to 68% water, 3% to 30% edible oil having an SFI of at least 25 to 70°F, 4% to 15% non-fat milk solids, and minor but effective amounts of stabilizer and emulsifier to form a mix;

20

homogenizing said mix to form an emulsion of fat globules having a particle size wherein the average  $d_{vs}$  value is 0.2 to 1.5 microns and a particle size distribution wherein the ratio of  $d_{vs}$  value is 0.2 to 1.5 microns and a particle size distribution wherein the ratio of  $d_{vs}$  to  $d_{max}$  has a quotient of from 9 to 14;

25

30

aging the homogenized mix at 30°F to 45°F for 2 to 12 hours to crystallize said fat globules;

35



whipping the aged mix under freezing conditions  
to an overrun of 50% to 110%; and

5 substantially retaining said particle size and  
particle size distribution after said  
whipping and freezing.

7. A process as defined in claim 6, characterized in  
10 that the oil and emulsifier are melted together and  
injected just prior to homogenization into a solution of  
the remaining ingredients in the water.

15 8. A process as defined in claim 6, characterized in  
that the aerated frozen dessert is hardened at -40°F to  
-80°F after whipping.

20 9. A process as defined in claim 8, characterized in  
that the hardened aerated frozen dessert is stored,  
distributed, sold and consumed at 0°F to 10°F.

---

25 10. A process as defined in claim 8, characterized in  
that 2 to 3 parts of the aerated frozen dessert are  
diluted at room temperature with one part of liquid and  
stirred to form a shake beverage.

30 11. A process for preparing a stable shake beverage,  
characterized in that the process comprises:

5 mixing 63% to 68% water, 3% to 9% edible oil  
having an SFI of at least 25 to 70°F, 4% to  
15% non-fat milk solids, and minor but  
effective amounts of stabilizer and  
emulsifier to form a mix;

10 homogenizing said mix to form an emulsion of fat  
globules having a particle size wherein the  
average  $d_{vs}$  value is 0.2 to 1.5 microns and  
a particle size distribution wherein the  
ratio of  $d_{max}$  to  $d_{vs}$  has a quotient of from  
9 to 14;

15 aging the homogenized mix at 30°F to 45°F for 2  
to 12 hours to crystallize said fat  
globules;

20 whipping the aged mix under freezing conditions  
to an overrun of 50% to 70%;

substantially retaining said particle size and  
particle size distribution after said  
whipping and freezing;

25 placing the aerated frozen product into  
individual containers; and

30 allowing said aerated frozen product to  
equilibrate to a temperature of from 15°F  
to 25°F until it develops a shake  
consistency.

12. A process as defined in claim 11, characterized in  
35 that the aerated frozen product is hardened at -40°F to  
-80°F after it is placed into individual containers.

13. A stable aerated frozen food product characterized in that it is formed by a process which comprises the steps of:

5        mixing 45% to 68% water, 3% to 30% edible oil  
         having an SFI of at least 25 to 70°F, 4% to  
         15% non-fat milk solids, and minor but  
         effective amounts of stabilizer and  
         emulsifier to form a mix;

10       homogenizing said mix to form an emulsion of fat  
         globules having a particle size wherein the  
         average  $d_{vs}$  value is 0.2 to 1.5 microns and  
         a particle size distribution wherein the  
15       ratio  $d_{vs}$  to  $d_{max}$  has a quotient of from 9  
         to 14;

         aging the homogenized mix at 30°F to 45°F for 2  
         to 12 hours to crystallize said fat  
20       globules;

         whipping the aged mix under freezing conditions  
         to an overrun of 50% to 110%; and

25       substantially retaining said particle size and  
         particle size distribution after said  
         whipping and freezing.

30 14. A stable ready to serve shake beverage, characterized  
     in that it is formed by a process which comprises:

- 5 mixing 63% to 68% water, 3% to 9% edible oil  
having an SFI of at least 25 to 70°F, 4% to  
15% non-fat milk solids, and minor but  
effective amounts of stabilizer and  
emulsifier to form a mix;
- 10 homogenizing said mix to form an emulsion of fat  
globules having a particle size wherein the  
average  $d_{vs}$  value is 0.2 to 1.5 microns and  
a particle size distribution wherein the  
ratio of  $d_{max}$  to  $d_{vs}$  has a quotient of from  
9 to 14;
- 15 aging the homogenized mix at 30°F to 45°F for 2  
to 12 hours to crystallize said fat  
globules;
- 20 whipping the aged mix under freezing conditions  
to an overrun of 50% to 70%;
- substantially retaining said particle size and  
particle size distribution after said  
whipping and freezing;
- 25 placing the aerated frozen product into  
individual containers; and
- 30 allowing said aerated frozen product to  
equilibriate to a temperature of from 15°F  
to 25°F until it develops a shake  
consistency.

15. A product as defined in claim 14, characterized in  
35 that it has an overrun of 50% to 110%.

0147433

16. A product as defined in claim 14 or claim 15,  
characterized in that at least 94% of said oil is  
dispersed in said aqueous phase in the form of said  
crystallized fat particles and less than 6% of said oil is  
5 in the form of de-emulsified fat.

17. A food product whenever formed by the process as  
claimed in any one of claims 1 to 12.  
10

18. A stable aerated frozen food product, characterized  
in that it comprises:

15 45% to 68% water in the form of ice crystals and  
an aqueous phase;

4% to 15% non-fat milk solids;

20 3% to 30% edible oil having an SFI of at least  
25 at 70°F, a majority of said oil being  
dispersed in said aqueous phase in the form  
of crystallized fat particles having a  $d_{vs}$   
value of 0.2 to 1.5 microns and a ratio of  
25  $d_{max}$  to  $d_{vs}$  having a quotient of from 9 to  
14; and

minor, but effective, amounts of emulsifier and  
stabilizer.

30

19. A product as defined in claim 18, characterized in  
that it has an overrun of 50% to 110%.

35

20. A product as defined in claim 18, characterized in that at least 94% of said oil is dispersed in said aqueous phase in the form of said crystallized fat particles and less than 6% of said oil is in the form of de-emulsified fat.

21. A product as defined in claim 18, characterized in that it further comprises a hydrocolloid stabilizer system in a quantity less than 1%.

22. A product as defined in claim 18, characterized in that it comprises 0.1% to 0.5% emulsifier.

23. A stable frozen mellorine product, characterized in that it comprises:

52% to 57% water in the form of ice crystals and an aqueous phase;

4% to 15% non-fat milk solids;

10% to 15% edible oil having an SFI of at least 25 at 70°F, at least 94% of said oil being dispersed in said aqueous phase in the form of crystallized fat particles having a  $d_{vs}$  value of 0.2 to 0.5 microns and a ratio of  $d_{max}$  to  $d_{vs}$  having a quotient of from 9 to 14, and less than 6% of said oil in the form of de-emulsified fat;

18% to 33% added sugars; and  
an overrun of 60% to 90%.

5  
24. A stable ready-to-serve shake beverage characterized  
in that it comprises:

10 63% to 68% water in the form of ice crystals and  
an aqueous phase:

4% to 15% non-fat milk solids;

15 3% to 9% edible oil having an SFI of at least 25  
at 70°F, at least 94% of said oil being  
dispersed in said aqueous phase in the form  
of crystallized fat particles having  $ad_{vs}$   
value of 0.2 to 1.5 microns and a ratio of  
20  $d_{max}$  to  $d_{vs}$  having aquotient of from 9 to  
14, and less than 6% of said oil in the  
form of de-emulsified fat;

18% to 33% added sugars;  
25 0.1% to 0.3% mono- and/or di-glyceride emulsifiers;  
less than 0.3% of a hydrocolloid stabilizer; and  
an overrun of 50% to 70%.



European Patent  
Office

# EUROPEAN SEARCH REPORT

**0147483**  
Application number

EP 83 11 3183

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 3)
Y	US-A-3 914 440 (RALPH P. WITZIG) * Column 2, lines 5-14; column 3, line 61 - column 4, line 3 *	1,2,6, 9,17	A 23 G 9/02 A 23 G 9/04
Y	US-A-4 421 778 (MARVIN L. KAHN et al.) * Column 3, lines 25-27; column 6, lines 25-30; claims 1,25 *	1,2-6, 9,10, 17	
Y	GB-A-2 045 050 (D. TRESSER) * Page 3, lines 50-60; page 2, lines 20-27 *	1,2,4- 6,17	
Y	US-A-4 368 211 (J.R. BLAKE et al.) * Column 7, lines 20-25; examples 1-3; column 8, lines 30-41 *	1-10, 17	TECHNICAL FIELDS SEARCHED (Int. Cl. 3)
Y	MILK PLANT MONTHLY, June 1948, pages 42-50; L.L. LITTLE: "Emulsifying and stabilizing agents for ice cream" * Page 44, column 2, paragraph 2; page 50, column 2, paragraph 4 *	1	A 23 G
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 27-07-1984	Examiner GUYON R.H.
<p><b>CATEGORY OF CITED DOCUMENTS</b></p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons &amp; : member of the same patent family, corresponding document</p>			





European Patent  
Office

# EUROPEAN SEARCH REPORT

0147483

Application number

EP 83 11 3183

Page 2

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl. 7)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
Y	US-A-4 031 261 (J.R. DURST)  * Examples 1-13; column 3, lines 14-24; claim 15; column 12, lines 30-35; column 6, lines 50-68 *	1, 9, 10, 17	
A	US-A-3 431 117 (G.J. LORANT) * Column 2, lines 34-37; column 3, lines 45-50; column 3, lines 15-17; claims 1-8 *	1	
A	US-A-3 889 001 (N.F. BUIDE et al.)		
A	FR-A- 868 427 (SOC. ASTRA) * Abstract no. 1a *	7	
A	US-A-4 346 120 (R.G. MORLEY et al.)		
A	US-A-3 949 102 (J.A. HELLYER et al.)		TECHNICAL FIELDS SEARCHED (Int. Cl. 7)
E, X	US-A-4 434 186 (NITIN-DESIA et al.)	1-24	
A	US-A-3 993 793 (D.J. FINNEY)		
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 27-07-1984	Examiner GUYON R.H.
<b>CATEGORY OF CITED DOCUMENTS</b>			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons  & : member of the same patent family, corresponding document	

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**